J Abnorm Child Psychol (2014) 42:311–320
DOI 10.1007/s10802-013-9775-3

Assessing Social Anxiety in African American Youth using the Social Phobia and Anxiety Inventory for Children

Armando A. Pina · Michelle Little · Henry Wynne · Deborah C. Beidel

Published online: 20 July 2013
© Springer Science+Business Media New York 2013

Social anxiety disorder is characterized by a persistent fear of social or performance situations in one or more areas, including public speaking, dating, and/or talking to new or unfamiliar people. Typically with an age of onset in adolescence, social anxiety is accompanied by evaluation concerns, functional impairment, and is prospectively and concurrently linked to substance use, un-employment, and dependence on the welfare system (Lipsitz and Schneier 2000; Morris et al. 2005; Tolman et al. 2009). Whereas there is ample literature about social anxiety in children and adolescents, most research has been based on White samples (Hunter and Schmidt 2010; Neal and Turner 1991). Turning to multiethnic samples, cross-ethnic comparative research based on rating scales shows that in clinical samples African American youth in particular report significantly lower (or similar) social anxiety levels than their Caucasian counterparts (e.g., Beidel et al. 2000, 1999; Ferrell et al. 2004). On average, African Americans could truly be low (or same as Caucasians) on social anxiety, but it also might be the case that measures are no adequately capturing social anxiety in African American youth. As reviewed by Pina et al. (2013), several measures have failed to provide equivalent information across ethnic groups, including for African American youth. Configural invariance in a community sample of White and African American youth, for example, was not supported for the Child Behavior Checklist (internalizing/externalizing scales; Tyson et al. 2011). Moreover, among African Americans anxiety seems to manifest itself largely in terms of physical symptoms (Neal and Turner 1991) and measures such as the Social Anxiety Scale for Adolescents (La Greca and Lopez 1998) that do not include physiological anxiety items might be under-identifying socially anxious African American youth. Clearly, the implication is that lack of invariance can result in poor science, overpathologizing, and wasted resources. As such, it is important to investigate whether measures developed with White samples provide equivalent information about ethnic minority youth in general, including for African Americans.

When it comes to cross-ethnic measurement invariance, item response theory and sophisticated quantitative methods can offer rich information about a scale's performance, especially compared to simple reliability analyses. Briefly, configural invariance tests yield information on whether the same factors of a measure exist across groups (Ghorpade et al. 1999; Millsap and Yun-Tein 2004; Vandenberg and Lance 2000), weak invariance can elucidate whether the items of a scale have the same meaning across groups (Labouvie and Ruetsch 1995; Raykov 2004), strong invariance tests can offer insights about the level or severity of anxiety needed for respondents to endorse a given item on a scale (Widamen and Reise 1997), and strict invariance refers to the error or unexplained variance in the endorsement of an item (Byrne et al. 1989). Armed with these methods and to shed some light on the usefulness of social anxiety measures

Author Note This study was supported in part by Award Numbers K01MH086687 and L60MD001839 from the National Institute of Mental Health and the National Center on Minority Health and Health Disparities. The content of this article is solely the responsibility of the authors and does not necessarily represent the official views of the funding agencies. We would like to thank Roger Millsap, Department of Psychology, Arizona State University, Daniel Sass, Department of Educational Psychology, University of Texas San Antonio and Augustine Osman, Department of Psychology, University of Texas San Antonio for their methodological consultation.

A. A. Pina (✉) · H. Wynne
Department of Psychology, Arizona State University, P.O. Box 871104, Tempe, AZ 85287, USA
e-mail: armando.pina@asu.edu

M. Little
Psychology Department, University of Texas at San Antonio, San Antonio, TX 78209, USA

D. C. Beidel
Department of Psychology, University of Central Florida, Orlando, FL 32816, USA

for the assessment of African American youth, we conducted secondary data analyses corresponding to a convenience sample of African American and White youth who completed the Social Phobia and Anxiety Inventory for Children (SPAI-C; Beidel et al. 1995). We focused on the SPAI-C because it is the most widely used social anxiety self-rating scale and it has been found to be sensitive to change in intervention trials (e.g., Beidel et al. 2007; Masia-Warner et al. 2005) thereby making it a measure with clinical utility. In addition, the SPAI-C does contain physiological anxiety items (Beidel et al. 1996, 1995) making it a good initial target for investigating, in a preliminary way, the assessment of social anxiety in African American youth. As such, the main goal of this study was to explore the measurement invariance and optimal cut-off scores of the SPAI-C for African American youth. To achieve this aim, primary analyses focused on using exploratory structural equation modeling to determine factor structure and test the cross-ethnic measurement invariance of the SPAI-C (see Asparouhov and Muthén 2009; Millsap 2011). In addition, Receiver Operator Characteristic analyses were used to examine optimal cut-off scores for the cross-ethnic group predictive validity of the SPAI-C. Optimal cutoff scores also were ascertained based on Youden's index (Youden 1950).

Method

Participants

Data corresponding to a convenience sample of 501 youth (8 to 16 years, M_{age}=11.62 years, *SD*=2.6, 249 girls; 120 African American, 381 Caucasian) were examined in this study. Cross-ethnic comparisons along age, sex and income revealed significant differences between African American and White youth in terms of income (lower income for African Americans, χ^2=7.53, p=0.023 Hollingshead Classification System; Hollingshead and Redlich 1958) and age (African Americans were slightly older M_{age}=12.25, *SD*=2.53, but less than one year on average t=−3.07, p=0.002; for White M_{age}=11.43, *SD*=2.58). About 24 % of African Americans were upper class, 44 % (n=24) middle class, and 32 % (n=17) lower class; among Whites, 33 % (n=50) were upper class, 52 % (n=79) middle class, and 15 % (n=22) lower class.

The sex by race chi-square was not significant ($\chi 2$=0.40, p=0.53). Based on the Anxiety Disorders Interview Schedule for Children (ADIS-IV: C/P; Silverman and Albano 1996), about 49 % of the sample met criteria for a primary social anxiety disorder diagnosis, 8 % met criteria for other anxiety disorders as the primary diagnosis and 4 % met criteria for other non-anxiety disorder diagnosis. In addition, about 40 % of the sample did not meet criteria for a diagnosis and were considered "typically developing youth" also based on the ADIS: C/P. For African Americans, 49 % met criteria for social anxiety disorder, 2 % met criteria other disorders, and 49 % were typically developing youth. For Whites, 49 % met criteria for social anxiety disorder, 9 % met criteria other disorders, and 37 % were typically developing youth. In general, there were significantly more African American youth in the typically developing group compared to Caucasians [$\chi 2$=13.66, p=0.003].

Measures

The *Social Phobia and Anxiety Inventory for Children (SPAI-C;* Beidel et al. 1995*)* is comprised of 26 items reflecting potentially fearful social situations. For each item, youth are given three choices from which they select the one that best describes how they feel, think, and behave in the situation. Items are scored as 0 ("never, or hardly ever"), 1 ("sometimes") or 2 ("most of the time, or always" rated) and the total scale score is used to derive clinical cutoffs. In Beidel et al. (1995), SPAI-C scores were significantly correlated with youth's self-rated trait anxiety (r=0.50) and fear levels (r=0.53). Moreover, a two-week retest reliability estimate of 0.86 and an alpha coefficient of 0.95 were found for the SPAI-C in Beidel et al. (1995).

The *Anxiety Disorders Interview Schedule for DSM-IV: Child and Parent Versions* (ADIS: C/P; Silverman and Albano 1996) is a semi-structured diagnostic interview focusing on anxiety and related disorders. The child and parent versions were administered to children and parents, respectively. The ADIS-C/P's manual describes administration procedures and process for deriving diagnoses, including comorbid diagnoses (Albano and Silverman 1996). The ADIS-IV: C/P yields reliable anxiety symptom counts (ICCs 0.78 to 0.95 for ADSI-C; 0.81 to 0.96 for ADIS-P), diagnoses (kappas 0.80 to 0.92), and clinical severity ratings (ADIS-CSR; rs 0.80 to 0.84) (Silverman et al. 2001). The ADIS-IV: C/P was the primary anxiety measure used to derive diagnoses in this sample.

Procedures

Participants for this study were recruited from the community (e.g., through referrals from pediatricians, social workers, and psychologists). After parents signed consent and youth provided assent, youth completed the SPAI-C as part of a comprehensive assessment battery that included the ADIS-IV: C/P (Silverman and Albano 1996). All study procedures were approved by the university's Institutional Review Board.

Data Analytic Plan

Exploratory structural equation modeling (ESEM) was used to determine factor structure and test measurement invariance

of the SPAI-C across ethnicity/race (and also sex). Sex was tested in the preliminary analyses because girls have been found to typically report greater anxiety levels than boys (Lewinsohn et al. 1998; Muris and Broeren 2009; Strauss and Last 1993), although this difference is not typically found in clinical samples. Given that our focus was on a clinic referred sample, sex was tested to render the ethnicity/race analyses more robust. We used an ESEM approach because it offers added precision in nested model invariance testing by reducing sources of model misfit in both large and small sample conditions (Asparouhov and Muthén 2009). Further, since SPAI-C items have a 3-point response scale, a weighted least squares mean variance (WLSMV) estimator was used, which is robust to violations of normality (Flora and Curran 2004; Muthén and Muthén 1998–2011). Invariance tests began with a nested multi-group "omnibus test" of the cross-group equality of the indicator covariance and mean structure matrices, which provides preliminary evidence for measurement non-invariance when significant (Millsap 2011). Next, configural analyses examined the overall model fit and significance of hypothesized factor loadings for a multi-group model with no cross-group constraints to ensure that the same factor structure was supported across groups.[1] In accord with Muthén's recommendations, strong invariance was tested next by comparing a model with cross-group factor loadings and item thresholds equality constraints to the configural model (Muthén and Asparouhov 2002; Widamen and Reise 1997). This strong invariance model provided a simultaneous test of the equivalence of the magnitude of item factor loadings and thresholds across groups of interest; thus implying cross-group equivalence of item meaning and item severity with respect to the anxiety construct. Further, strict invariance was tested by comparing a model with constrained loadings, thresholds, and item residuals to a model with constrained loadings and thresholds but free item residuals. Strict invariance comparisons assess equivalence of cross-group item consistency. Finally, factor structure differences across groups were compared using factor variance-covariance and latent mean equality constraints in separate sets of nested model tests.

Model fit for full sample and configural models was evaluated on the basis of the chi-square measure of absolute fit and two practical fit indices: the comparative fit index (CFI), and root mean square error of approximation (RMSEA). Cutoffs of CFI ≥ 0.95 and RMSEA ≤ 0.05 suggest good fit in baseline configural models and RMSEAs between 0.06 and 0.08 suggest adequate fit (Cheung and Rensvold 2002; Hu and Bentler 1998). Subsequent measurement non-invariance was evaluated on the basis of a majority of indices including (1) significant change in the chi-square between successive nested models,[2] (2) change in the RMSEA of 0.007 or more and (3) a change in the CFI of ≤ -0.002 (Meade et al. 2008; Sass et al. in press). Chi-square change and practical fit cutoffs were selected to optimize power and minimize Type I error for detection of measurement non-invariance using WLSMV estimation with our small samples (Elosua 2011; Meade et al. 2008; Sass et al. in press).

Receiver Operator Characteristic (ROC) analyses were used to examine optimal cut-off scores for the predictive validity of the SPAI-C across sex and race. Cross-sex assessment of optimal cut-offs was evaluated given that sometimes girls report higher anxiety symptom levels (Muris and Broeren 2009), and to provide a useful comparison point for interpretation of cross-race ROC analyses. ROC analyses examined SPAI-C prediction of (i) any DSM IV anxiety disorder diagnoses for diagnosed and typically developing youth and (ii) any social anxiety disorder diagnosis for social phobic and typically developing youth. From ROC analyses, area under the curve (AUC) was derived by plotting the sensitivity (SN) of SPAI-C cutoff scores against the false positive rate [1-specificity (SP)] of SPAI-C cutoff scores ranging from 1 to 50. Note that SN yields the proportion of diagnosed individuals detected at a given SPAI-C score while SP yields the proportion of undiagnosed individuals correctly identified at a given SPAI-C score. AUCs in the 0.80 to 0.90 range are considered "good" whereas AUCs in the 0.90 to 1.00 range are considered "excellent". Points on the ROC curve indicating a balance between optimally high SN and low false positive rates (1-SP) represent optimal cutoff scores for disease detection. Youden's index [(SN + SP)-1] was used to evaluate the balance between SN and SP. Higher Youden's Index scores indicate better cutoffs (Youden 1950).

Results

Preliminary Analyses

ESEM with an oblique rotation was used to evaluate the fit and factor solutions of full sample models ranging from 3 to 5 factors because prior research supports SPAI-C factor solutions of 3 and 5 factors (Beidel 1996; Beidel et al. 1995). Table 1 summarizes model fit and interpretability of the factor solutions. A 4-factor solution was selected as optimal given that it showed an overall model fit in the good

[1] The ESEM configural model was identified in accord with Muthén's recommendations for ESEM multi-group models with categorical variables and theta parameterization (Muthén and Muthén 1998–2011). Specifically, all item residuals were initially constrained to 1 in both groups and latent factor means were constrained to 0 in both groups. Further, to estimate the ESEM model across groups, latent factor variances were set to 1 across groups. Once factor loadings and thresholds were constrained to equality across groups for the strong invariance tests, constraints on latent factor means and variances and item residuals were released in the second non-reference group.

[2] Chi-square differences using the WLSMV estimator were ascertained from an algorithm based on Asparouhov et al. (2006).

Table 1 Overall ESEM model fit and interpretability of 3–5 factor solutions of the SPAI-C

	3-factor	4-factor	5-factor
χ^2 Fit (*df*)	350.852*** (99)	250.764*** (107)	202.616*** (102)
CFI	0.956	0.975	0.982
RMSEA	0.072	0.053	0.045
Factor Interpretability	3 factors interpretable	4 factors interpretable	4 factors interpretable

***$p<0.001$

range [χ^2 (107, N=479)=250.764, $p<0.001$; CFI=0.975, RMSEA=0.053] and four fully interpretable factors that matched factors previously identified in published SPAI-C factor analyses (see Table 2; Beidel 1996; Beidel et al. 1995). In contrast, RMSEA fit of a 3-factor model was just adequate [χ^2 (99, N=479)=350.852, $p<0.001$; CFI=0.956, RMSEA=0.072] and the 5-factor model revealed only 4 interpretable factors. As shown in Table 2, two factors of the 4-factor solution matched corresponding Assertiveness and General Conversation and Public Performance factors identified previously with the exception that item 8 of the current Public Performance factor formerly loaded on the Assertiveness and General Conversation factor (Beidel et al. 1995). The two remaining factors also matched Avoidance and Physical and Cognitive Symptoms factors identified in a prior 5-factor solution, although item 22 of the current Physical and Cognitive Symptoms and item 7 of the current Avoidance factor were not represented in the prior 5-factor solution (Beidel 1996). The selected 4-factor loading structure was used as a target structure in subsequent cross-group ESEM models.

Cross-Sex Measurement and Factor Invariance of the SPAI-C

Preliminary omnibus test of equality of the covariance and mean structure matrices suggested potential cross-sex measurement non-invariance. Specifically, constraining item covariances and thresholds to equality across sex resulted in a significant chi-square change and a decrement in practical fit [$\Delta\chi^2$ (26, N=479)=44.64, $p<0.05$; ΔCFI=−0.004; ΔRMSEA=0.05]. Next, a cross-sex configural ESEM model showed that the full sample targeted 4-factor structure was not replicated adequately. Specifically, three items (1, 7 and 23) did not show highest and adequate loadings (i.e. significant, standard loadings > 0.30) on target factors. Re-testing the cross-sex configural model with those items removed showed adequate fit [χ^2 (128, N=479)=248.233, $p<0.001$; CFI=0.976; RMSEA=0.062] and a cross-sex loading pattern resembling the target factor pattern (see Table 3). All items showed their largest, significant factor loadings on corresponding target factors.

Subsequent tests supported cross-sex measurement invariance of the SPAI-C, but highlighted latent mean differences. As shown in Table 4, constraining factor loading and thresholds to equality across sex resulted in a significant chi-square change, but practical fit was not adversely affected [$\Delta\chi^2$ (59, N=479)=89.034, $p<0.01$; ΔCFI=0.003; ΔRMSEA=−0.007], thus supporting cross-sex strong measurement invariance. Follow-up strict invariance tests showed negligible change in model fit indices, therefore affirming cross-sex invariance of item residuals. Similarly, invariance of factor variances and covariances was affirmed when factor variances and covariances were constrained to equality across sex. Finally, cross-sex factor mean differences were revealed by significant decrements in chi-square, CFI and RMSEA fit when latent means were constrained [$\Delta\chi^2$ (2, N=479)=24.809, $p<0.001$; ΔCFI=−0.004; ΔRMSEA=0.023]. Specific latent mean differences were tested by regressing latent means on gender in a full sample model.[3] Compared to girls, boys showed lower Public Performance (b=−0.37, $p<0.001$), Assertiveness and General Conversation (b=−0.44, $p<0.001$) and Physical and Cognitive Symptoms (b=−0.22, $p<0.05$).

Cross-Ethnic Measurement and Factor Invariance of the SPAI-C

Preliminary omnibus test of the equality of the covariance and mean structure matrices supported cross-ethnic measurement invariance of the SPAI-C. Constraining item covariances and thresholds to cross-ethnic equality resulted in negligible change in chi-square and CFI tests, although the RMSEA worsened [$\Delta\chi^2$ (23, N=479)=26.94, $p>0.05$; ΔCFI=0.001; ΔRMSEA=0.027]. A subsequent cross-race 4-factor configural model using all items revealed that 3 items (1, 9 and 14) did not show adequate loadings on target factors across race. As shown in Table 4, with items 1, 9 and 14 removed from the configural model, the resulting model fit was good [χ^2 (118, N=479)=193.958, $p<0.001$; CFI=0.984, RMSEA=0.051]. All factor loadings reflected the target structure (see Table 5). Cross-ethnic measurement invariance of the SPAI-C was affirmed by measurement and factor invariance tests (see Table 4). Constraints on loadings and thresholds and then item residuals had negligible effects on chi-square and practical fit. Similarly, constraining latent factor variances/covariances and latent means did not affect chi-square or practical fit.

[3] Partial invariance tests of latent factor means is not possible in ESEM.

Table 2 Standardized factor loadings of the 4-factor ESEM SPAI-C model

No. Item	Factors			
	Public performance	Assertiveness general conversation	Avoidance	Physical cognitive symptoms
Public performance				
1 scared when joining a large group	**0.428**		0.335	
2 scared when becoming the center of attention	**0.404**			
3 scared when I have to do something while others watch me	**0.687**			
4 scared when speaking or reading in front of a group	**0.88**			
5 scared when answering questions in class or at group meetings	**0.677**			
8 too scared to ask questions in class	**0.577**			
16 scared when speaking in front of the class	**0.843**			
17 scared when in a school play, choir, music, or dance recital	**0.697**			
Assertiveness & general conversation				
9 scared in the school cafeteria		**0.375**		
10 scared if someone starts arguing		**0.665**		
11 scared if someone asks me to do something that I don't want to do		**0.65**		
12 scared in an embarrassing situation		**0.561**		0.327
13 scared if someone says something that is wrong or bad		**0.697**		
14 scared when I start to talk to someone	0.319	**0.544**	0.32	
15 scared if I have to talk for longer than a few minutes	0.304	**0.454**		
18 scared when ignored or made fun of by others		**0.572**		
Avoidance				
6 scared at parties, dances, school, and go home early			**0.557**	
7 scared to meet new kids			**0.392**	
19 I avoid social situations (parties, school, playing with others)			**0.702**	
20 I leave social situations			**0.746**	
23 I don't speak until spoken to			**0.388**	
Physical & cognitive symptoms				
21 before going to a party, I think about what might go wrong				**0.615**
22 I am unable to speak or sound funny when talking to others				**0.563**
24 when I am with other people, I think "scary" thoughts				**0.593**
25 before going someplace, I feel (somatic symptoms)				**0.906**
26 when I am in a social situation, I feel (somatic symptoms)				**0.898**

All represented factor loadings are significant at $p<0.05$. Bolded factor loadings represent primary factor loadings for particular factors. Factor loadings that are not bolded are non-zero cross-loadings above 0.30. Item 8 formerly factored on the Assertiveness/General Conversation Factor (Beidel et al. 1995). Items 7 and 22 did not load on any factors in prior research (Beidel 1996; Beidel et al. 1995)

Sensitivity, Specificity, and Clinical Meaningfulness of the SPAI-C

We next examined sensitivity, specificity, and clinical meaningfulness analyses across sex and race. Table 6 shows results from Receiver Operator Characteristic (ROC) analyses, including AUC, sensitivity, and specificity for models predicting (i) a social anxiety disorder diagnosis in a sub-sample of social phobic and typically developing youth and (ii) any DSM IV anxiety disorder diagnosis in a sub-sample sample of diagnosed (86 % had social anxiety disorder based on the ADIS-C/P) and typically developing youth. Figure 1a–d depict corresponding ROC curves for both African-American and Caucasian groups. In the prediction of any anxiety disorder, Youden's index suggested a cutoff of 14 as the criterion with the best combination of SN and SP for girls and White youth (range 0.79 to 0.86). For African American youth, however, Youden's index suggested a cutoff of 17 was best (SN/SP=0.74/0.90), although SN was below 0.80. Focusing on optimizing SN, however, a cut-off of 15 provides high SN and minimal loss in SP for African-American youth (SN/SP=0.81/0.83). Similarly, although Youden's index indicated an optimal cut-off of 14 for boys, an alternative cut-off of 13 shows higher SN (0.78) and acceptable SP (0.79) for boys.

Table 3 Cross-sex standard factor loadings of the 4-factor configural ESEM SPAI-C model. ESEM item cross loadings are not shown in the Table

Item #	Public performance		Assertiveness & general conversation		Avoidance		Physical & cognitive symptoms	
	Boys	Girls	Boys	Girls	Boys	Girls	Boys	Girls
2	0.329	0.594						
3	0.786	0.742						
4	0.973	0.825						
5	0.882	0.591						
8	0.662	0.481						
16	0.764	0.900						
17	0.561	0.867						
9			0.391	0.642				
10			0.709	0.873				
11			0.772	0.850				
12			0.699	0.709				
13			0.850	0.872				
14			0.627	0.867				
15			0.481	0.801				
18			0.667	0.692				
6					0.357	0.604		
19					0.436	0.811		
20					0.740	0.865		
21							0.706	0.548
22							0.572	0.628
24							0.728	0.502
25							0.952	0.813
26							0.908	0.839

All represented loadings are significant at $p<0.05$

In the prediction of a social anxiety disorder diagnosis specifically, Youden's index suggested 14 as an optimal cutoff for boys, girls, and White youth. For African-American youth, a cutoff of 17 was suggested by Youden's index, which yielded a SN of 0.75 and a SP of 0.90. Focusing on optimal SN/SP, however, a cut-off of 15 provides higher SN and minimal loss in SP for African-American youth (SN/SP=0.82/0.83).

Table 4 Summary of model fit statistics for measurement and factor structure invariance tests

	$\chi^2(df)$	CFI	RMSEA	$\Delta\chi2(df)$	ΔCFI	ΔRMSEA
Cross-sex						
Configural invariance	248.233*** (128)	0.976	0.062			
Strong invariance				89.034** (59)	0.003	−0.007
Strict invariance				25.505 (18)	0.006	−0.002
Factor variance/Covariances invariance				1.267 (3)	0.014	−0.015
Latent means invariance				24.809***(2)	−0.004	0.023
Cross-race						
Configural invariance	193.958*** (118)	0.984	0.051			
Strong invariance				55.242 (52)	0.006	−0.010
Strict invariance				26.784 (17)	0.000	0.002
Factor variance/Covariances invariance				4.150 (3)	0.008	−0.004
Latent means invariance				2.063 (2)	0.000	−0.001

Factor loadings and thresholds are constrained for strong invariance tests. Item residuals, loadings and thresholds are constrained for strict invariance tests

** $p<0.01$, *** $p<0.001$

Table 5 Cross-race standard factor loadings of the 4-factor configural ESEM SPAI-C model

Item #	Public performance		Assertiveness & general conversation		Avoidance		Physical & cognitive symptoms	
	White	African American	White	African American	White	African American	White	African American
2	0.354	0.557						
3	0.716	0.633						
4	0.896	0.877						
5	0.684	0.590						
8	0.584	0.555						
16	0.803	0.957						
17	0.650	0.837						
10			0.691	0.792				
11			0.654	0.815				
12			0.635	0.586				
13			0.733	0.927				
15			0.418	0.463				
18			0.523	0.883				
6					0.537	0.757		
7					0.421	0.516		
19					0.671	0.920		
20					0.732	0.789		
23					0.357	0.371		
21							0.634	0.643
22							0.606	0.694
24							0.622	0.508
25							0.968	0.847
26							0.970	0.791

ESEM item cross loadings are not shown in the Table. All represented loadings are significant at $p<0.05$.

Table 6 Receiver Operator Characteristics Curve (ROC) Analyses of the SPAI-C: differences by gender, ethnicity, and optimal cutoff

	AUC	Confidence Interval		Cutoff of 18		Cutoff of 15		Cutoff of 14		Cutoff of 13	
		LC	UC	SN	SP	SN	SP	SN	SP	SN	SP
Model 1 Any anxiety disorder vs. Typically developing youth											
Full Sample	0.887	0.857	0.917	0.70	0.90	0.79	0.83	0.81	0.82	0.83	0.79
Afr.Am	0.905	0.851	0.958	0.71	0.92	0.81	0.83	0.81	0.81	0.83	0.79
Caucasian	0.882	0.845	0.918	0.70	0.89	0.78	0.83	0.81	0.83	0.83	0.79
Female	0.889	0.847	0.931	0.76	0.87	0.85	0.80	0.86	0.79	0.87	0.77
Male	0.881	0.838	0.923	0.64	0.92	0.72	0.86	0.75	0.85	0.78	0.79
Model 2 Social anxiety disorders vs. Typically developing youth											
Full Sample	0.891	0.860	0.922	0.72	0.90	0.80	0.83	0.83	0.82	0.87	0.74
Afr.Am	0.909	0.856	0.963	0.71	0.92	0.82	0.83	0.82	0.81	0.84	0.79
Caucasian	0.885	0.848	0.923	0.72	0.89	0.80	0.83	0.83	0.83	0.84	0.79
Female	0.899	0.856	0.942	0.78	0.87	0.87	0.80	0.88	0.79	0.88	0.77
Male	0.886	0.842	0.930	0.65	0.93	0.74	0.86	0.77	0.85	0.79	0.81

AUC is area under the curve. *SN* is sensitivity. *SP* is specificity. *Afr. Am.* is African-American/Black. All AUCs are significant at the $p < 0.01$ level



Fig. 1 **a** ROC curve of SPAI-C prediction of any anxiety disorder among Caucasians **b** ROC curve of SPAI-C prediction of any anxiety disorder among African-Americans **c** ROC curve of SPAI-C prediction of social anxiety disorder among Caucasians **d** ROC curve of SPAI-C prediction of social anxiety disorder among African-Americans

Discussion

Findings from the present study suggest that social anxiety levels among African American youth based on the SPAI-C appear to be "true" and not a function of measurement bias. In addition, since measurement invariance for the SPAI-C was largely ascertained in this study, findings support past research showing that African American youth generally have significantly lower (or similar) social anxiety levels than their White counterparts (e.g., Beidel et al. 2000, 1999; Ferrell et al. 2004).

In the present study, cross-ethnic non-invariance for a handful of SPAI-C items was found. Even if these findings replicate, non-invariance for a few items is not necessarily a cause for concern (Widamen and Reise 1997). Knowledge about the invariant items: "scared when joining a large group", "scared in the school cafeteria", and "scared when I talk to someone" can help refine the cultural sensitivity of the SPAI-C. In particular, it has been suggested that socially anxious African American youth show more anxiety in situations with same-race youth compared to situations with Whites (Neal and Ward-Brown 1994). Neal-Barnett and Smith (1997) explain that African Americans sometimes fear being accused of "Acting White" when meeting same race-individuals. In addition, we believe it is possible for African American youth to interpret these items (e.g., *scared when joining a large group*) in the context of predominantly 'White situations' which might activate for some youth racial socialization survival skills rather than anxiety. As such, exploring the various meanings African American youth might be assigning to non-invariant items can inform any SPAI-C content revision and its cultural sensitivity in the assessment of African American youth.

Turning to cutoff scores, our results showed that the previously suggested SPAI-C cutoff of 18 might be under-identifying anxious youth. Instead, lower SPAI-C cutoff scores (in the 13 to 15 range) appear to be more adequate. These lower cutoffs have implications for screening youth into prevention efforts, identifying more cases in need of diagnostic "work-ups" and treatment services, and for better estimating program effects (Aune and Stiles 2009; Beidel et al. 2007; Masia-Warner et al. 2005). Lower cutoffs can help address health disparities among African American youth because they have been found to be less likely to use specialty care without identification or encouragement (Alegría et al. 2012). As such, these revised scores can facilitate casting a wider net to serve more African American youth and families.

The present study findings have clinical practice implications, including for working in the contexts of cultural diversity. Clinicians using the SPAI-C in their practice should specifically question African American youth who endorse any of the identified non-invariant items. This questioning should include asking the child to generate examples that pertain to the specific item. In terms of clinical cutoffs, we suggest cautiously using the scores suggested herein (less than the traditional 18) with the caveat that revised scores are not likely to distinguish youth with social anxiety disorder from youth with other anxiety disorders. This lack of

disorder specificity within the anxieties is commonly linked to symptom overlap (Silverman and Ollendick 2005). It also is important to note that our cross-sex invariance analyses yielded non-invariant items and very similar revised cut-offs for boys versus girls. We feel it is difficult to untangle whether sex differences are (un)related to ethnicity since our sample size prohibited use of 4-way multiple group analyses (smallest group would have been less than 70). As such, asking African American youth about endorsed invariant items should not be culturally-guided, but open to the possibility that gender may play a role.

Several limitations and directions for future research are noteworthy. First, data used for the present study was drawn from a convenience sample of families who sought services for the child's anxiety. In the case of African American families, youth were likely identified as anxious and parents encouraged to seek help (Alegría et al. 2012). For these reason, findings might not be completely representative of African American youth and any nuanced ways they experience anxiety (Kingery et al. 2007). Second, as noted earlier, our sample size prohibited use of 4-way multiple group analyses to untangle any possible sex-by-race/ethnicity effects. This is an important future step given our findings that boys endorsed less public performance, assertiveness and general conversation symptoms while a few of the non-invariant items identified for African Americans resided in those same factor scales. Third, whereas our focus on African American youth was sample driven, other cultural groups with diverse views and interpretations of mental illness should be a focus of these types of investigations as well (Knight et al. 2009) since cultural socialization relevant to symptom interpretation and expression can bias clinical assessment results when using measures developed for another cultural group (see Pina et al. 2013). As data accumulate to identify evidence-based assessments that are culturally robust, greater progress can be made to improve the mental health of ethnic minority youth and families.

References

Albano, A. M., & Silverman, W. K. (1996). *Guide to the use of the anxiety disorders interview schedule for DSM-IV-child and parent versions*. London: Oxford University Press.

Alegría, M., Lin, J. Y., Green, J. G., Sampson, N. A., Gruber, M. J., & Kessler, R. C. (2012). Role of referrals in mental health service disparities for racial and ethnic minority youth. *Journal of the American Academy of Child and Adolescent Psychiatry, 51*, 703–711. doi:10.1016/j.jaac.2012.05.005.

Asparouhov, T., & Muthén, B. (2009). Exploratory structural equation modeling. *Structural Equation Modeling: A Multidisciplinary Journal, 16*, 397–438. doi:10.1080/10705510903008204.

Asparouhov, T., Muthén, L. Muthén, B. O. (2006). *Robust chi square difference testing with mean and variance adjusted test statistics*. Retrieved from http://www.statmodel.com/download/webnotes/webnote10.pdf.

Aune, T., & Stiles, T. C. (2009). Universal-based prevention of syndromal and subsyndromal social anxiety: a randomized controlled study. *Journal of Consulting and Clinical Psychology, 77*, 867–879. doi:10.1037/a0015813.

Beidel, D. C. (1996). Assessment of childhood social phobia: construct, convergent, and discriminative validity of the Social Phobia and Anxiety inventory for Children (SPA-C). *Psychological Assessment, 8*, 235. doi:10.1037/1040-3590.8.3.235.

Beidel, D. C., Turner, S. M., & Morris, T. L. (1995). A new inventory to assess childhood social anxiety and phobia: the social phobia and anxiety inventory for children. *Psychological Assessment, 7*, 73–79. doi:10.1037/1040-3590.7.1.73.

Beidel, D. C., Fink, C. M., & Turner, S. M. (1996). Stability of anxious symptomatology in children. *Journal of Abnormal Child Psychology, 24*, 257–269. doi:10.1007/BF01441631.

Beidel, D. C., Turner, S. M., & Morris, T. L. (1999). Psychopathology of childhood social phobia. *Journal of the American Academy of Child and Adolescent Psychiatry, 38*, 643–650. doi:10.1097/00004583-199906000-00010.

Beidel, D. C., Turner, S. M., Hamlin, K., & Morris, T. L. (2000). The Social Phobia and Anxiety Inventory for Children (SPAI-C): external and discriminative validity. *Behavior Therapy, 31*, 75–87. doi:10.1016/S0005-7894(00)80005-2.

Beidel, D. C., Turner, S. M., Sallee, F. R., Ammerman, R. T., Crosby, L. A., & Pathak, S. (2007). SET-C versus fluoxetine in the treatment of childhood social phobia. *Journal of the American Academy of Child and Adolescent Psychiatry, 46*, 1622–1632. doi:10.1097/chi.0b013e318154bb57.

Byrne, B. M., Shavelson, R. J., & Muthén, B. (1989). Testing for the equivalence of factor covariance and mean structures: the issue of partial measurement invariance. *Psychological Bulletin, 105*, 456–466. doi:10.1037/0033-2909.105.3.456.

Cheung, G. W., & Rensvold, R. B. (2002). Evaluating goodness-of-fit indexes for testing measurement invariance. *Structural Equation Modeling, 9*, 233–255. doi:10.1207/S15328007SEM0902_5.

Elosua, P. (2011). Assessing measurement equivalence in ordered-categorical data. *Psicológica, 32*, 403–421. Retrieved from http://www.uv.es/psicologica/articulos2.11/13ELOSUA.pdf.

Ferrell, C. B., Beidel, D. C., & Turner, S. M. (2004). Assessment and treatment of socially phobic children: a cross cultural comparison. *Journal of Clinical Child and Adolescent Psychology, 33*, 260–268. doi:10.1207/s15374424jccp3302_6.

Flora, D. B., & Curran, P. J. (2004). An empirical evaluation of alternative methods of estimation for confirmatory factor analysis with ordinal data. *Psychological Methods, 9*, 466–491. doi:10.1037/1082-989X.9.4.466.

Ghorpade, J., Hattrup, K., & Lackritz, J. R. (1999). The use of personality measures in cross-cultural research: a test of three personality scales across two countries. *Journal of Applied Psychology, 84*, 670–679. doi:10.1037/0021-9010.84.5.670.

Hollingshead, A. B., & Redlich, F. C. (1958). *Social class and mental illness: Community study*. New York: John Wiley.

Hu, L., & Bentler, P. (1998). Fit indices in covariance structure modeling: sensitivity to under parameterized model misspecification. *Psychological Methods, 3*, 424–453.

Hunter, L. R., & Schmidt, N. B. (2010). Anxiety psychopathology in African American adults: literature review and development of an empirically informed sociocultural model. *Psychological Bulletin, 136*, 211–235. doi:10.1037/a0018133.

Kingery, J., Ginsburg, G. S., & Alfano, C. A. (2007). Somatic symptoms and anxiety among African American adolescents. *Journal of Black Psychology, 33*, 363–378. doi:10.1177/0095798407307041.

Knight, G. P., Roosa, M. W., & Umaña-Taylor, A. J. (2009). Measurement and measurement equivalence issues. In G. P. Knight, M. W.

Roosa, & A. J. Umaña-Taylor (Eds.), *Studying ethnic minority and economically disadvantaged populations: Methodological challenges and best practices* (pp. 97–134). Washington, DC: American Psychological Association. doi:10.1037/11887-004.

La Greca, A. M., & Lopez, N. (1998). Social anxiety among adolescents: linkages with peer relations and friendships. *Journal of Abnormal Child Psychology, 26*, 83–94. doi:10.1023/A:1022684520514.

Labouvie, E., & Ruetsch, C. (1995). Testing for equivalence of measurement scales: simple structure and metric invariance reconsidered. *Multivariate Behavioral Research, 30*, 63–76. Retrieved from http://login.ezproxy1.lib.asu.edu/login?url=http://search.proquest.com/docview/618682253?accountid=4485.

Lewinsohn, P. M., Gotlib, I. H., Lewinsohn, M., Seeley, J. R., & Allen, N. B. (1998). Gender differences in anxiety disorders and anxiety symptoms in adolescents. *Journal of Abnormal Psychology, 107*, 109–117. doi:10.1037/0021-843X.107.1.109.

Lipsitz, J. D., & Schneier, F. R. (2000). Social phobia: epidemiology and cost of illness. *PharmacoEconomics, 18*, 23–32. doi:10.2165/00019053-200018010-00003.

Masia-Warner, C., Klein, R. G., Dent, H. C., Fisher, P. H., Alvir, J., Albano, A. M., et al. (2005). School-based intervention for adolescents with social anxiety disorder: results of a controlled study. *Journal of Abnormal Child Psychology, 33*, 707–722. doi:10.1007/s10802-005-7649-z.

Meade, A. W., Johnson, E. C., & Braddy, P. W. (2008). Power and sensitivity of alternative fit indices in tests of measurement invariance. *Journal of Applied Psychology, 93*, 568. doi:10.1037/0021-9010.93.3.568.

Millsap, R. E. (2011). *Statistical approaches to measurement invariance*. New York: Routledge.

Millsap, R. E., & Yun-Tein, J. (2004). Assessing factorial invariance in ordered-categorical measures. *Multivariate Behavioral Research, 39*, 479–515. doi:10.1207/S15327906MBR3903_4.

Morris, E. P., Stewart, S. H., & Ham, L. S. (2005). The relationship between social anxiety disorder and alcohol use disorders: a critical review. *Clinical Psychology Review, 25*, 734–760. doi:10.1016/j.cpr.2005.05.004.

Muris, P., & Broeren, S. (2009). Twenty-five years of research on childhood anxiety disorders: publication trends between 1982 and 2006 and a selective review of the literature. *Journal of Child and Family Studies, 18*, 388–395. doi:10.1007/s10826-008-9242-x.

Muthén, B., & Asparouhov, T. (2002). *Latent variable analysis with categorical outcomes: Multiple-group and growth modeling in Mplus.* Retrieved from http://statmodel2.com/download/webnotes/CatMGLong.pdf.

Muthén, L. K., & Muthén, B. O. (1998-2011). *Mplus user's guide* (6th ed.). Los Angeles: Muthén & Muthén.

Neal, A. M., & Turner, S. M. (1991). Anxiety disorders research with African Americans: current status. *Psychological Bulletin, 109*, 400–410. doi:10.1037//0033-2909.109.3.400.

Neal, A. M., & Ward-Brown, B. J. (1994). Fears and anxiety disorders in African American children. In S. Friedman (Ed.), *Anxiety disorders in African Americans* (pp. 65–75). New York: Springer.

Neal-Barnett, A., & Smith, J. (1997). African Americans. In S. Friedman (Ed.), *Cultural issues in the treatment of Anxiety* (pp. 154–174). New York: Guilford Press.

Pina, A. A., Gonzales, N., Holly, L. E., Zerr, A., & Wynne, H. (2013). Toward evidence-based assessment of ethnic minority youth. In B. D. McLeod, A. Jensen-Doss, & T. Ollendick (Eds.), *Diagnostic and behavioral assessment in children and adolescents*. New York: Guilford.

Raykov, T. (2004). Behavioral scale reliability and measurement invariance evaluation using latent variable modeling. *Behavior Therapy, 35*, 299–331. doi:10.1016/S0005-7894(04)80041-8.

Sass, D. A., Schmitt, T. A., & Marsh, H. W. (in press). Evaluating model fit with ordered categorical data within a measurement invariance framework: a comparison of estimators. *Structural Equation Modeling*.

Silverman, W. K., & Albano, A. M. (1996). *Anxiety disorders interview schedule for children*. San Antonio: Psychological Corporation.

Silverman, W. K., & Ollendick, T. H. (2005). Evidence-based assessment of anxiety and its disorders in children and adolescents. *Journal of Clinical Child and Adolescent Psychology, 34*, 380–411. doi:10.1207/s15374424jccp3403_2.

Silverman, W. K., Saavedra, L. M., & Pina, A. A. (2001). Test-retest reliability of anxiety symptoms and diagnoses with anxiety disorders interview schedule for DSM-IV: child and parent versions. *Journal of the American Academy of Child and Adolescent Psychiatry, 40*, 937–944. doi:10.1097/00004583-200108000-00016.

Strauss, C. C., & Last, C. G. (1993). Social and simple phobias in children. *Journal of Anxiety Disorders, 7*, 141–152. doi:10.1016/0887-6185(93)90012-A.

Tolman, R. M., Himle, J., Bybee, D., Abelson, J. L., Hoffman, J., & Van Etten-Lee, M. (2009). Impact of social anxiety disorder on employment among women receiving welfare benefits. *Psychiatric Services, 60*, 61–66. doi:10.1176/appi.ps.60.1.61.

Tyson, E. H., Teasley, M., & Ryan, S. (2011). Using the child behavior checklist with African American and Caucasian American adopted youth. *Journal of Emotional and Behavioral Disorders, 19*, 17–26. doi:10.1177/1063426609357760.

Vandenberg, R. J., & Lance, C. E. (2000). A review and synthesis of the measurement invariance literature: suggestions, practices, and recommendations for organizational research. *Organizational Research Methods, 3*, 4–69. Retrieved from http://login.ezproxy1.lib.asu.edu/login?url=http://search.proquest.com/docview/619672309?accountid=4485.

Widamen, K. F., & Reise, S. P. (1997). Exploring the measurement invariance of psychological instruments: Applications in the substance abuse domain. In K. J. Bryant (Ed.), *Alcohol and substance abuse research* (pp. 281–324). Washington, D.C.: American Psychological Association.

Youden, W. J. (1950). Index for rating diagnostic tests. *Cancer, 3*, 32–35. doi:10.1002/1097-0142(1950)3:1<32::AID-CNCR2820030106>3.0.CO;2-3.